

Hypercomplex Seminar 2023

July 9–14

Program and abstracts

(2023-07-10)

Program	1	(→)
Abstracts	7	(→)

July 9, 2023

Opening ceremony (17:45-17:55)

Session (Inaugural lectures),
Chairmen: Dariusz Partyka and Mariusz Zubert

- 18:00-18:50** *Sergiy Plaksa*, (→)
Monogenic functions in infinite-dimensional vector spaces with a commutative multiplication and harmonic vectors
- 19:00-19:50** *Marek Danielewski*, (→)
Quaternion Quantum Mechanics II: Unraveling the Mysteries of Gravity and the Dirac Equation within the Planck-Kleinert Crystal
- 20:00-20:50** *Chantal Roth*, (→)
Quaternionic foundations of Quantum Mechanics and spin 1/2 visualization

July 10, 2023

Session A1 (Hypercomplex structures),
Chairman: Sergiy Plaksa

- 10:00-10:50** *Lino F. Reséndis O.*, (→)
 \mathcal{Q}_p spaces for hyperholomorphic functions in the unit ball of \mathbb{R}^4
- 11:00-11:25** *Serhii Gryshchuk*, (→)
Representations of solutions of Lamé system with real coefficients via monogenic functions in the biharmonic algebra
- 11:30-12:20** *Massimo Vaccaro*, (→)
The full set of invariants characterizing the $Sp(n)$ -orbits in the real Grassmannians $G^{\mathbb{R}}(k, 4n)$

Session A2 (Complex and real analysis of one variable),
Chairman: Lino F. Reséndis Ocampo

- 13:00-13:25** *Mariia Stefanchuk*, (→)
On the exponential asymptotic of solutions of the nonlinear Cauchy-Riemann-Beltrami type equation

July 11, 2023

Session A3 (Holomorphic functions of one variable), Chairman: Serhii Gryshchuk

- 10:00-10:25** *Iryna Denega*, (→)
Evolutionary type inequalities for products of inner radii
- 10:30-10:55** *Maciej Parol*, (→)
The Koebe radius for certain class of polynomials
- 11:00-11:25** *Szymon Ignaciuk*, (→)
Physical interpretation of the membership to Kaplan classes of certain functions
- 11:30-11:55** *Jacek Dziok*, (→)
On multivalent prestarlike functions

Session A4 (Complex and real analysis of several variables), Chairman: Massimo Vaccaro

- 12:30-12:55** *Valery Volchkov, Vitaly Volchkov*, (→)
Continuous mean periodic extension of functions from a segment
- 13:00-13:25** *Anna Kimaczyńska*, (→)
The grad div operator

Session A5 (Poster session), Chairman: Massimo Vaccaro

- 13:30-14:00** *Olga Rovenska*, (→)
Approximation of classes of Poisson integrals by Fejér means
- 13:30-14:00** *Natalia Zorii*, (→)
On the role of the point at infinity in Deny's principle of positivity of mass for Riesz potentials

July 12, 2023

Session B1 (Physics: Ontology of Quantum Mechanics), Chairman: Krzysztof Pomorski

- 10:00-10:50** *Jarosław Duda, (→)*
Topological charges with electromagnetic + gravitomagnetic interactions
- 11:00-11:50** *Richard Kerner, (→)*
Non-linear electrodynamics derived from the Kaluza-Klein 5-dimensional gravity
- 12:30-13:20** *Manfried Faber, (→)*
Running coupling from a classical soliton model
- 13:30-13:55** *Łukasz Stępień, (→)*
On some exact solutions of certain nonlinear partial differential equations
- 14:00-14:50** *Jarosław Duda, (→)*
Maximal Entropy Random Walk to understand Born rule and Bell violation

Session A6 & B (Advanced mathematical methods in physics), Chairman: Jarosław Duda

- 15:00-15:50** *Osamo Suzuki, (→)*
Binary and ternary structures in evolution theory (VI) An entropic description of the evolution of the universe
- 16:00-16:25** *Małgorzata Nowak, (→)*
Iterative construction of discrete Laplacians on 2D square and hexagonal lattices using BTBB and other sequences

July 13, 2023

Session C1 & B (Quantum Computer Science),
Chairman: Chantal Roth

- 10:00-10:50** *K. Gnatenko, (→)*
Geometric measure of entanglement of variational quantum states and its quantifying on a quantum computer
- 11:00-11:50** *K. Pomorski, D. Kotula, (→)*
Classical and Quantum Conway Game of Life

Session C2 & B (Classical/Quantum Electronics and Machine Learning),
Chairman: Khrystyna Gnatenko

- 12:30-12:55** *M. Kowalik, (→)*
Machine Learning in Cryogenic Experiments
- 13:00-13:25** *M. Zubert, Z. Kulesza, M. Jankowski, A. Napieralski, (→)*
Application of Scattering Parameters to The DPL Time-Lag Parameter Estimation using MEMS Structures

Session D1 & B (Quantum Electronics),
Chairman: Mariusz Zubert

- 13:30-13:55** *M. Złotkowski, (→)*
Local systems management of different energy sources

Discussion panel (14:00-14:30)
Chairmen: Sergiy Plaksa, Manfred Faber, Chantal Roth

July 14, 2023

Session D2 & B (Quantum Electronics), Chairman: Krzysztof Pomorski

- 10:00-10:50** *J. Sosnowski*, (→)
Influence of quantized vortex dynamics on superconducting electronics
- 11:00-11:25** *V. Vachtsevanos*, (→)
Emergence of topological defects in finite 2-D lattices during second phase order transitions
- 11:30-12:20** *M. Belogolosvskii*, (→)
Two new issues in superconducting electronics: transparent samples and SQUID-like Josephson junctions
- 12:30-12:55** *Oleh Yermakov*, (→)
Discovery of polarization degree of freedom for localized light

Session A7 (Geometrical methods in physics), Chairman: Sergiy Plaksa

- 13:30-13:55** *A. Paszkiewicz*, (→)
Does the convex Peano curve exist and what can we say about operators?
- 14:00-14:25** *Armen Grigoryan*, (→)
Landau and Bloch theorems for planar harmonic functions

Session C3 & B & D (Computer Science in Quantum Electronics), Chairman: Manfred Faber

- 14:30-14:55** *K. Pomorski*, (→)
Theory of interface between semiconductor quantum dots and Josephson junction
- 15:00-15:50** *B. Stojewski, K. Pomorski*, (→)
Hybrid Schroedinger-GL algorithm in solving problems
- 16:00-16:25** *Ł. Pluszyński, K. Pomorski*, (→)
Solving Schroedinger equation with analog electronics
- 16:30-17:55** *Ł. Pluszyński*, (→)
Simulation and experiments with superconducting single photon detectors

Closing ceremony (18:00-18:15)

July 15, 2023

Jarosław Duda and Krzysztof Pomorski
recommend topics on Ontology of Quantum Mechanics
(QM Foundations & Nature of Time seminar)

16:00-17:00 *Sergey Rashkovskiy (RAS)*, <http://th.if.uj.edu.pl/~dudaj/QMNoT>
Quantum Mechanics: Strange Particle Theory or Classical Field Theory?

MIKHAIL BELOGOLOVSKII

Department of Experimental Physics, Faculty of Mathematics, Physics and Informatics,
Comenius University
(Bratislava, Slovak Republic)
(belogolovskii@ukr.net)

Two new issues in superconducting electronics: transparent samples and SQUID-like Josephson junctions

Abstract. Nowadays, superconducting circuits are among the most prominent platforms for quantum computing, communications, and sensing. Transparent superconductors are of particular interest, since they allow direct interfacing between stationary superconducting and flying photonic qubits. The latter are necessary for the transmission of quantum information because their quantum states are more resistant to decoherence. I present a methodological approach for the development of superconductors with acceptable transparency and usefully high critical temperature, followed by discussion of representative characteristics of a number of already known oxides. In particular, I examine how the complex interplay of free carrier absorption and interband transitions can lead to the coexistence of comparatively high carrier density at the Fermi level and the needed transparency for visible light. To date, noticeable successes have been achieved using the first approach and, as a good example of its implementation, we overview detailed results obtained for electrochemically doped indium-tin-oxide compounds, which is a reference material for testing our ideas.

Next, I present our results on synthetic binary superlattices formed by nanometer-thick layers of normal and ferromagnetic metals, quantum transport through which unexpectedly demonstrates the formation of edge states, which are resistant to disorder and other mobile charge interactions. The idea of the experiments was inspired by a paradigmatic explanation of the plateaus in transversal transport characteristics in a 2D conductor at very low temperatures and strong magnetic fields (the integer quantum Hall effect) that is based on the existence of narrow near-boundary quantum channels of non-interacting electrons created at the Fermi level in strongly disordered electron systems. The edge states are chiral in the sense that they can carry current only in one fixed direction. At the same time, the edge modes are topologically protected and their number cannot vary under continuous transformation of the system. This approach assumes the presence of a disordered and isotropic two-dimensional electron gas. Our aim has been to create a strongly anisotropic 3D system with edge channels concentrated mainly at the hinges of the structure, i.e., a synthetic material with one or two 1D conducting modes. One of the best ways to probe spatial current distributions is Josephson interferometry in a stacked configuration where the hybrid structure under study links two superconducting (S) electrodes. The figure of merit in such experiments is the maximum supercurrent (I_c) versus in-plane magnetic field (H). In our experiments, we applied this technique to study the transport of Cooper pairs through a periodic multilayer formed by ten normal metal (Al) - ferromagnetic metal (Ni or Ni-Fe alloy) bilayers. For the first time, we observed SQUID-like ($I_c - H$) oscillations instead of conventional Fraunhofer patterns expected for trivial S-weak link-S junctions. Possible applications of such devices are discussed.

Keywords and phrases: Josephson junctions, transparent superconducting films, ferromagnetic-normal metallic superlattices, Josephson interferometry

(←)

MAREK DANIELEWSKI
 Faculty of Applied Mathematics, AGH UST (Kraków, Poland)
 (daniel@agh.edu.pl)

Quaternion Quantum Mechanics II: Unraveling the Mysteries of Gravity and the Dirac Equation within the Planck-Kleinert Crystal

Abstract. We present quaternion representation of quantum mechanics that allows its ontological interpretation. The correspondence between classical and quaternion quantum equations, permits considering the universe (vacuum) as an ideal elastic solid. Elementary particles would have to be standing or soliton-like waves. Tension induced by the compression and twisting of the elastic medium would increase the energy density, consequently generate a gravity forcing and affect the wave speed. Consequently the gravity could be described by an index of refraction.

Theory was created by combining the Cauchy model of the elastic continuum with the Planck-Kleinert crystal hypothesis. The quaternion-imaginary Lagrangian, the quaternion motion equation and the quaternionic oscillator allowed deriving:

- The Schrödinger equation from the functional integral, which identifies the quaternion-imaginary quantum Hamiltonian.
- The 2nd order wave equation system describing both the bosons and the gravity.
- The first order quaternionic wave equation system.
- The family of the second order wave equation systems describing both the particles and the generated quaternionic force-fields.
- The quaternionic continuity equation in an ideal elastic solid.

Keywords and phrases: quaternion

Joint work with: Lucjan Sapa, Chantal Roth

(←)

IRYNA DENEGA

Institute of mathematics of the National Academy of Sciences of Ukraine (Kyiv, Ukraine)
(iradenega@gmail.com)

Evolutionary type inequalities for products of inner radii

Abstract. In this presentation we consider evolutionary type inequalities for the following products

$$\begin{aligned} & r(B_0, 0) \prod_{k=1}^n r(B_k, a_k), \\ & r(B_\infty, \infty) \prod_{k=1}^n r(B_k, a_k), \\ & r(B_0, 0) r(B_\infty, \infty) \prod_{k=1}^n r(B_k, a_k), \end{aligned}$$

where $n \in \mathbb{N}$, $A_n = \{a_k\}_{k=1}^n$ be an arbitrary fixed system of points of the complex plane, B_0 , B_∞ and $\{B_k\}_{k=1}^n$ be an arbitrary system of domains such that $a_0 = 0 \in B_0 \subset \overline{\mathbb{C}}$, $\infty \in B_\infty \subset \overline{\mathbb{C}}$, $a_k \in B_k \subset \overline{\mathbb{C}}$, $k = \overline{1, n}$, $B_i \cap B_j = \emptyset$, $i \neq j$, $r(B, a)$ denote an inner radius of the domain B with respect to a point a .

Keywords and phrases: the Green function, an inner radius of the domain, logarithmic capacity

REFERENCES

- [1] A. K. Bakhtin, I. V. Denega, *Generalized M.A. Lavrentiev's inequality*, J. Math. Sci. **262** (2022), 138–153, (<https://doi.org/10.1007/s10958-022-05806-y>).
- [2] I. Denega, *Extremal decomposition of a multidimensional complex space with poles on the boundary of a polydisk*, In: Cerejeiras P., Reissig M., Sabadini I., Toft J. (eds) Current Trends in Analysis, its Applications and Computation. Trends in Mathematics. Birkhäuser, Cham. (2022), 143–151, (https://doi.org/10.1007/978-3-030-87502-2_14).
- [3] A. K. Bakhtin, I. V. Denega, *Weakened problem on extremal decomposition of the complex plane*, Matematychni Studii **51** (2019), 35–40, (<https://doi.org/10.15330/ms.51.1.35-40>).

(←)

JAROSŁAW DUDA

Institute of Computer Science, Jagiellonian University (Kraków, Poland)
(jaroslaw.duda@uj.edu.pl)

Maximal Entropy Random Walk to understand Born rule and Bell violation

Abstract. Turns out standard random walk/diffusion usually only approximates the (Jaynes) maximal entropy principle - necessary for statistical mechanics models. MERW (maximal entropy random walk) as e.g. random walk along Ising sequence repairs this approximation, this way also disagreements with quantum mechanics - like Anderson localization, Born rule and Bell violation. I will introduce to MERW and discuss suggestions for physics understanding, including potential new possibilities and their applications.

Slides here: <https://www.dropbox.com/s/a8yqfabq3gxsjth/Bell%20mini.pdf>

(←)

JAROSŁAW DUDA

Institute of Computer Science, Jagiellonian University (Kraków, Poland)
(jaroslaw.duda@uj.edu.pl)

Topological charges with electromagnetic + gravitomagnetic interactions

Abstract. In liquid crystals they experimentally obtain topological charges with long-range e.g. Coulomb-like interactions. It brings question how far can we take its resemblance with particle physics. I will talk about such looking promising approach based on Landau-de Gennes model, with EM-like Lagrangian interpreting field curvature as dual EM field (Faber's approach). This way Gauss law counts topological charge for its quantization, also thanks to Higgs-like potential regularizing charge to finite energy in agreement with the running coupling effect. Then there appear further particle-like topological defects, like topological vortices with knots resembling baryons (e.g. having proton lighter than neutron), and nuclei knotted against Coulomb repulsion. Extending to 4D field adding boosts, their dynamics turns out governed by second set of Maxwell equations for gravity.

Slides here: <https://www.dropbox.com/s/9d12g91ypzqu5hp/liquid%20crystal%20particles.pdf>

(←)

JACEK DZIOK
University of Rzeszów (Rzeszów, Poland)
(jdzio@ur.edu.pl)

On multivalent prestarlike functions

Let \mathcal{A}_p ($p \in \mathbb{N} := \{1, 2, \dots\}$) denote the class of functions of the form

$$f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n$$

which are analytic in $\mathcal{U} := \{z \in \mathbb{C} : |z| < 1\}$. A function $f \in \mathcal{A}_p$ is said to be *multivalent starlike of order α* ($0 \leq \alpha < p$) if

$$\operatorname{Re} \left(\frac{z f'(z)}{f(z)} \right) > \alpha \quad (z \in \mathcal{U}).$$

We denote by $\mathcal{S}_p^c(\alpha)$ the class of all functions which are starlike of order α .

Due to Ruscheweyh [1] we introduce the class of multivalent prestarlike functions. We say that a function $f \in \mathcal{A}_p$ belongs to the class $\mathcal{R}_p(\alpha)$ of *multivalent prestarlike functions of order α* if

$$f(z) * \frac{z^p}{(1-z)^{2(p-\alpha)}} \in \mathcal{S}_p^*(\alpha),$$

where "*" denotes the Hadamard product (or convolution).

The main object of the talk is to present some properties of multivalent prestarlike functions and their applications.

REFERENCES

- [1] S. Ruscheweyh, *Convolutions in geometric function theory*, Sem. Math. Sup. 83, Les Presses de l'Université de Montréal, 1982.

(←)

MANFRIED FABER
Technische Universität (Wien, Austria)
(faber@kph.tuwien.ac.at)

Running coupling from a classical soliton model

Abstract. Running coupling in field theory was found by the necessity to cancel the infinities which appear in the Coulomb field of point-like charges and experimentally verified in high energy collisions of electron-positron pairs. The model of topological particles (MTP) describes electrons as topological solitons of finite size and mass, of a field of rotations of spatial Dreibeins without any singularities. Therefore, there is no need to remove any infinities. The question whether MTP is able to describe the running of the electric charge is investigated in numerical computations of the energy of soliton pairs at varying distance. It is shown that within MTP the running charge of electrons originates naturally in the finite size of electrons.

(←)

KHRYSTYNA GNATENKO

Ivan Franko National University of Lviv and SoftServe Inc. (Lviv, Ukraine)

Geometric measure of entanglement of variational quantum states and its quantifying on a quantum computer

Abstract. The entanglement of variational quantum states is studied with analytical calculations and on quantum computers of IBM and Rigetti. We consider variational quantum states prepared with rotational RY, RZ, and RXX gates. The states have a graph structure and can be considered as quantum graph states constructed with the action of the operator of evolution corresponding to the Ising model. In the particular cases of parameters of RY and RZ gates the states were studied in [1]. The entanglement of the states is obtained analytically for the arbitrary structure of a graph. The dependencies of the entanglement on the parameters of the rotational gates are analyzed. The states are prepared on IBM's quantum computer. We calculate the geometric measure of entanglement of the states with quantum programming. The protocol for the calculations is based on the relation of the geometric measure of entanglement with mean spin obtained in [2].

We also study the entanglement of variational quantum states corresponding to the Generative Adversarial Network states [3, 4]. In the case of one-layer, the states also can be considered as quantum graph states [5]. The entanglement of the states is calculated analytically and with quantum programming on IBM's and Rigetti quantum computers [6, 7]. The results of quantum calculations are in good agreement with the theoretical ones.

Keywords and phrases: quantum computers, geometric measure of entanglement, quantum graph states, Generative Adversarial Network states

REFERENCES

- [1] Kh. P. Gnatenko, V. M. Tkachuk, *Entanglement of graph states of spin system with Ising interaction and its quantifying on IBM's quantum computer*, Phys. Lett. A. **396**, 127248, 5 p. (2021).
- [2] A. M. Frydryszak, M. I. Samar, V. M. Tkachuk
Quantifying geometric measure of entanglement by mean value of spin and spin correlations with application to physical systems Eur. Phys. J. D **71**, 233 [8 p.] (2017).
- [3] Kh. P. Gnatenko *Evaluation of variational quantum states entanglement on a quantum computer by the mean value of spin* arXiv:2301.03885 (2023).
- [4] Ch. Zoufal, Lucchi, A. Woerner, S. *Quantum Generative Adversarial Networks for learning and loading random distribution* npj Quantum Information **5**, 103 (2019).
- [5] Kh. P. Gnatenko, N. A. Susulovska *Geometric measure of entanglement of multi-qubit graph states and its detection on a quantum computer*, EPL (Europhys. Lett.) **136**, No. 4, 40003 (2021).
- [6] IBM Q experience. <https://quantum-computing.ibm.com/>
- [7] Rigetti Computing: Quantum Computing <https://www.rigetti.com/>

(←)

ARMEN GRIGORYAN

The John Paul II Catholic University of Lublin (Lublin, Poland)

(armen@kul.pl)

Landau and Bloch theorems for planar harmonic functions

Abstract. In this presentation we discuss a Landau type theorem for properly normalized bounded planar harmonic functions. We also present Bloch type theorems for properly normalized quasiregular harmonic functions and for open harmonic functions.

Keywords and phrases: planar harmonic functions, Landau theorem, Bloch theorem

(←)

SERHII GRYSHCHUK

Institute of Mathematics of NAS of Ukraine (Kyiv, Ukraine)
National University of Kyiv-Mohyla Academy (Kyiv, Ukraine)
(serhii.gryshchuk@gmail.com)

Representations of solutions of Lamé system with real coefficients via monogenic functions in the biharmonic algebra

Abstract.

Consider a Lamé (Lamé-type) system

$$(1) \quad \begin{cases} \Delta u(x, y) + p \frac{\partial \theta(x, y)}{\partial x} = 0, \\ \Delta v(x, y) + p \frac{\partial \theta(x, y)}{\partial y} = 0 \quad \forall (x, y) \in D, \end{cases}$$

where $\Delta := \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$, $p \in \mathbb{R} \setminus \{0\}$, $\theta := \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y}$.

By a solution of the Lamé system (1) we mean its *regular* solution, i.e., a pair of twice continuously differentiable functions (u, v) , which satisfies the equality (1). If $p = -1$ we assume that $\Delta u = \Delta v = 0$ additionally. For some positive values of p the system (1) is a Lamé system of equilibrium equations of isotropic plane theory of elasticity with respect to the vector of displacements $(u(x, y), v(x, y))$.

Among all two-dimensional algebras of the second rank with unity e over the field of complex numbers \mathbb{C} we found a *biharmonic* algebra $\mathbb{B} := \{c_1 e + c_2 \rho : c_k \in \mathbb{C}, k = 1, 2\}$, $\rho^2 = e$, containing bases $\{e_1, e_2\}$, such that a general solution of (1) can be rewritten with the help of components of \mathbb{B} -valued “analytic” functions $\Phi_k(xe_1 + ye_2)$ (x, y are real variables), $k = 1, 2$, and certain polynomials with the order less or equal to three.

REFERENCES

- [1] Gryshchuk S.V., *Commutative Complex Algebras of the Second Rank with Unity and Some Cases of Plane Orthotropy I*, Ukr. Math. J., Volume 70, Issue 8 (2019), pp. 1221-1236.
- [2] Gryshchuk S.V., *Commutative Complex Algebras of the Second Rank with Unity and Some Cases of Plane Orthotropy II*, Ukr. Math. J., Volume 70, Issue 10 (2019), pp. 1594-1603.
- [3] Gryshchuk S.V., *\mathbb{B}_0 -valued monogenic functions and their applications to the theory of anisotropic plane media*, In: *Analytic Methods of Analysis and Differential Equations: AMADE 2018* (Eds. S.V. Rogosin and M.V. Dubatovskaya), Cambridge: Cambridge Scientific Publishers Ltd, (2020), 33 – 48. ISBN (paperback): 978-1-908106-65-0.
- [4] Gryshchuk S.V., *Monogenic functions with values in commutative complex algebras of the second rank with unity and generalized biharmonic equation with simple nonzero simple characteristics*, Ukr. Math. Zh, Volume 73, Issue 4 (2021), pp. 474–487 (Ukrainian, English summary), DOI: 10.37863/umzh.v73i4.6199.

(←)

SZYMON IGNACIUK
University of Life Sciences in Lublin (Lublin, Poland)
(szymon.ignaciuk@up.lublin.pl)

Physical interpretation of the membership to Kaplan classes of certain functions

Abstract. The presented results concern a certain class of finite products of the form

$$(1) \quad \mathbb{D} \ni z \mapsto F_n(z; T_n; P_n) := \prod_{k=1}^n (1 - ze^{-it_k})^{p_k} ,$$

where $\mathbb{N} \ni n \mapsto T_n := (t_1, t_2, \dots, t_n)$ is an increasing sequence of values from $[0; 2\pi)$ such that $t_1 := 0$ and $\mathbb{N} \ni n \mapsto P_n := (p_1, p_2, \dots, p_n)$ is a sequence of real numbers of the same sign. The first results for polynomials with all zeros on unit circle \mathbb{T} (when P_n is a sequence of natural numbers) were given by Jahangiri [4] in terms of a gap condition. Complete membership to Kaplan classes for the polynomials was presented in [1]. In [2] the authors carried out complete membership to Kaplan classes of finite products of the form similar to (1), but with zeros symmetrically situated in \mathbb{T} . The presented results from [3] generalize ones from [1] and [2]. The interpretation of the obtained gap condition in terms of mass and density is given in the case where all zeros of the studied functions are situated in \mathbb{T} . An open problem is a more sophisticated physical interpretation that allows for the consideration of zeros outside the unit circle.

Keywords and phrases: Kaplan classes, critical points, mass, density

Joint work with: Maciej Parol

REFERENCES

- [1] S. Ignaciuk, M. Parol, *Zeros of complex polynomials and Kaplan classes*, Analysis Mathematica **46** (2020), 769–779. DOI: <https://doi.org/10.1007/s10476-020-0044-8>.
- [2] S. Ignaciuk, M. Parol, *Kaplan classes of a certain family of functions*, Annales UMCS Sectio A Mathematica **74(2)** (2020), 31–40. DOI: <https://doi.org/10.17951/a.2020.74.2.31-40>.
- [3] S. Ignaciuk, M. Parol, *A gap condition for the zeros of a certain class of finite products*, Proceedings of the Romanian Academy - Series A **22(1)** (2021), 19–25. [https://acad.ro/sectii2002/proceedings/doc2021-1/03-Ignaciuk Parol.pdf](https://acad.ro/sectii2002/proceedings/doc2021-1/03-Ignaciuk%20Parol.pdf)
- [4] M. Jahangiri, *A gap condition for the zeroes of certain polynomials in Kaplan classes $K(\alpha, \beta)$* , Mathematika **34** (1987), 53–63. DOI: <https://doi.org/10.1112/S0025579300013279>.

(←)

RICHARD KERNER

Laboratoire de Physique Théorique de la Matière Condensée, Sorbonne-Université, 4 Place Jussieu
(Paris, France)

(richard.kerner@sorbonne-universite.fr)

Non-linear electrodynamics derived from the Kaluza-Klein 5-dimensional gravity

Abstract. The lagrangian of the Kaluza-Klein theory, in its simplest five-dimensional version, should include not only the scalar curvature R , but also the quadratic Gauss-Bonnet invariant.

The metric tensor of the five-dimensional Kaluza-Klein space displays 15 degrees of freedom, including the space-time metric (10 components), the electromagnetic 4-potential (4 components) and the scalar field represented by the g_{55} component. We consider the case where only electromagnetic potential is present on the Minkowskian space-time, and in the absence of scalar field.

General lagrangian is computed and the resulting non-linear equations which generalize Maxwell's system in a quite unique way are investigated. A possibility of the existence of static solutions is presented, and the qualitative behaviour of such solutions as models for the electron is discussed.

(←)

ANNA KIMACZYŃSKA
 University of Lodz (Lodz, Poland)
 (anna.kimaczynska@wmii.uni.lodz.pl)

The grad div operator

Abstract. In this presentation we investigate some natural differential operators in the bundle of symmetric tensors on a Riemannian manifold M of dimension n . In particular we investigate two operators: the gradient and the divergence. The definition of these two operators is analogous to the definition of gradient and divergence in the bundle of skew-symmetric forms given by H. M. Rummeler [2]. It is interesting that the composition of the divergence and the gradient led to a second order linear differential operator div grad that was investigated in detail in the author's PhD dissertation [1]. However, the subject of our research here will be the grad div operator.

Keywords and phrases: Symmetric tensors, symmetric derivative, the gradient, the divergence.

REFERENCES

- [1] A. Kimaczyńska, *The differential operators in the bundle of symmetric tensors on a Riemannian manifold*, PhD thesis, Łódź University, Faculty of Mathematics and Computer Science, Łódź (2016), 59 pp;
- [2] H. M. Rummeler, *Differential forms, Weitzenböck formulae and foliations*, Publicacions Matemàtiques 33 (1989), 543–554.

(←)

KRZYSZTOF POMORSKI

Cracow Unversty of Technology-Faculty of Electrical and Computer Engineering, Quantum Hardware
(Kraków, Poland)

(kdvpomorski@gmail.com)

DARIUSZ KOTULA

Cracow Unversty of Technology-Faculty of Electrical and Computer Engineering, Quantum Hardware
(Kraków, Poland)

Different methodologies in description Classical and Quantum Conway Game of Life

Abstract. Classical stochastic Conway Game of Life is expressed by the dissipative Schrödinger equation and dissipative tight-binding model. This is conducted at the prize of usage of time dependent anomalous non-Hermitian Hamiltonians as with occurrence of complex value potential that do not preserve the normalization of wave-function and thus allows for mimicking creationism or annihilationism of cellular automaton. Simply saying time-dependent complex value eigenenergies are similar to complex values of resonant frequencies in electromagnetic resonant cavities reflecting presence of dissipation that reflects energy leaving the system or being pumped into the system. At the same time various aspects of thermodynamics were observed in cellular automata that can be later reformulated by quantum mechanical pictures. The usage of Shannon entropy and mass equivalence to energy points definition of cellular automata temperature. Contrary to intuitive statement the system dynamical equilibrium is always reflected by negative temperatures. Diffusion of mass, energy and temperature as well as phase of proposed wave function is reported and can be directly linked with second thermodynamics law approximately valid for the system, where neither mass nor energy is conserved. The concept of complex-valued mass mimics wave-function behavior. Equivalence an anomalous second Fick law and dissipative Schrödinger equation is given. Dissipative Conway Game of Life tight-binding Hamiltonian is given using phenomenological justification.

(←)

MAŁGORZATA NOWAK-KĘPCZYK
The John Paul II Catholic University of Lublin (Lublin, Poland)
(gosianmkk@wp.pl)

Iterative construction of discrete Laplacians on 2D square and hexagonal lattices using BTBB and other sequences

Abstract. The BTBB construction discussed in [1, 2, 3, 4] and its various applications were given in mathematics, biology, chemistry, knot theory, formal language theory. Several papers have been written on "Binary and ternary structures of the evolutions in the universe". Parts I-V by Suzuki, pointing out the uniqueness of this sequence.

We want to demonstrate iterative construction of discrete Laplacians on 2D square and hexagonal lattices using BTBB sequence and compare with the constructions obtained with the use of other sequences. The BTBB sequence occurs to be exceptional under many respects.

REFERENCES

- [1] J. Ławrynowicz, K. Nôno, and O. Suzuki, *Binary and ternary Clifford analysis vs. non-commutative Galois extensions. I. Basics of the comparison*, Bull. Soc. Sci. Lettres Łódź Sér. Rech. Déform. **62**, no. 1 (2012), 33–42.
- [2] J. Ławrynowicz, M. Nowak-Kępczyk, O. Suzuki, in collaboration with M. Othman, *Binary and ternary structures in the evolution of the universe III. The entropy description of the evolution*, Bull. Soc. Sci. Lettres Łódź Sér. Rech. Déform. **70** no. 1, (2020) 11–41. DOI: 10.26485/0459-6854/2020/70.1/1.
- [3] Julian Ławrynowicz, Małgorzata Nowak-Kępczyk, Osamu Suzuki, *Bi-nary and ternary structures of the evolutions in the universe (2x3x2x...-world) IV. The entropy description of evolution.*, Bull. Soc. Sci. Lettres Łódź Sér. Rech. Déform. **70** no. 1, (2020) 43–81. DOI: 10.26485/0459-6854/2020/70.1/2
- [4] M. Ławrynowicz, M. Nowak-Kępczyk, O. Suzuki, *Binary and ternary structures in the evolution of the universe V, Non-commutative Galois theory of evolution*, in: (HYPER)COMPLEX Seminar 2021 In Memoriam Of Prof. Julian Ławrynowicz. A Monograph, Soc. Sci. Lodziensis, Łódź 2021, pp. 7–37. DOI: 10.26485/978-83-60655-92-4/2

(←)

MACIEJ PAROL

The John Paul II Catholic University of Lublin (Lublin, Poland)

(mparol@kul.pl)

The Koebe radius for certain class of polynomials

Abstract. The presented results concern close-to-convex polynomials with all zeros of derivative in the unit circle \mathbb{T} . The minimal disc containing all images of the unit disc \mathbb{D} and the maximal disc contained in all images of \mathbb{D} through polynomials of degree 3 and 4 were determined in [4]. Moreover, all extremal functions for both problems were received. The same problem for polynomials of degree 5 was solved in [5]. In addition, the hypothesis for polynomials of odd degree was also given.

Joint work with: Szymon Ignaciuk

REFERENCES

- [1] Bieberbach L., *Über die Koeffizienten derjenigen Potenzreihen, welche eine schlichte Abbildung des Einheitskreises vermitteln*, Sitzungsber. Preuss. Akad. Wiss. Phys-Math. Kl., pp. 940-955, (1916).
- [2] Dmitrishin D., Dyakonov K., Stokolos A., *Univalent polynomials and Koebe's one-quarter theorem*, Anal. Math. Phys. **9**(3), pp. 991-1004, (2019). DOI: 10.1007/s13324-019-00305-x
- [3] Ignaciuk S., Parol M., *Zeros of complex polynomials and Kaplan classes*, Anal. Math. **46**, pp. 769-779, (2020). DOI: 10.1007/s10476-020-0044-8
- [4] Ignaciuk S., Parol M., *On the Koebe Quarter Theorem for certain polynomials*, Anal. Math. Phys. **11**(68), (2021). DOI: 10.1007/s13324-021-00501-8
- [5] Ignaciuk S., Parol M., *On the Koebe Quarter Theorem for certain polynomials of odd degree*, Anal. Math. Phys. **12**(92), (2022). DOI: 10.1007/s13324-022-00703-8

(←)

ADAM PASZKIEWICZ

University of Lodz, Faculty of Mathematics and Computer Science (Łódź, Poland)

(adam.paszkieicz@wmii.uni.lodz.pl)

Does the convex Peano curve exist and what can we say about operators?

Abstract. We present an unexpected construction of continuous surjection $f : [0, 1] \rightarrow T$ for a given closed compact set T in \mathbb{R}^2 . Some important consequences are connected with convexity preserving operators.

(\leftarrow)

SERGIY PLAKSA

Institute of Mathematics of the National Academy of Sciences of Ukraine (Kyiv, Ukraine)

University of Padova (Padova, Italy)

(plaksa62@gmail.com)

Monogenic functions in infinite-dimensional vector spaces with a commutative multiplication and harmonic vectors

Abstract. We consider special topological vector spaces with a commutative multiplication for some of elements of the spaces. The consideration of such topological vector spaces is motivated by the need to describe all spatial harmonic functions as components of hypercomplex monogenic functions. Monogenic functions are understood as continuous and differentiable in the sense of Gâteaux functions. We prove that all spatial harmonic functions are components of monogenic functions taking values in the mentioned spaces. We describe relations between the mentioned monogenic functions and harmonic vectors in the three-dimensional real space and establish sufficient conditions for infinite monogeneity of functions. Unlike the classical complex analysis, it is done in the case where the validity of the Cauchy integral formula for monogenic functions remains an open problem.

(←)

LINO F. RESÉNDIS O.

Universidad Autónoma Metropolitana-Azcapotzalco (Ciudad de México, México)

(lfro@azc.uam.mx)

\mathcal{Q}_p spaces for hyperholomorphic functions in the unit ball of \mathbb{R}^4

Abstract. The classical \mathcal{Q}_p for $1 \leq p < \infty$ spaces were introduced by R. Aulaskari and P. Lappan in [1] and for $0 < p \leq 1$ by R. Aulaskari et al in [2]. The hyperholomorphic generalization to the unit ball of \mathbb{R}^3 was given by K. Gürlebeck, et al in [3].

Consider the unit ball of $\mathbb{B} \subset \mathbb{R}^4$ and the set of real quaternions \mathbb{H} . For $0 \leq p < 2$, the $\tilde{\mathcal{Q}}_p(\mathbb{B})$ space consists of the \mathbb{H} right-module of hyperholomorphic functions $f : \mathbb{B} \rightarrow \mathbb{H}$ satisfying

$$\|f\| = \sup_{a \in \mathbb{B}} \int_{\mathbb{B}} |\overline{D}f(x)|^2 g^p(x, a) dx < \infty,$$

where $g(x, a) = \frac{1}{|\varphi_a(x)|^2} - 1$ is the composition of a modified fundamental solution of the Laplacian in the unit ball of \mathbb{B} , that is, $\frac{1}{|x|^2}$ and the Möbius transformation $\varphi_a(x) = (a - x)(1 - \bar{a}x)^{-1}$.

Let $0 \leq p < \infty$. The $\mathcal{Q}(\mathbb{B})$ space consists of the \mathbb{H} right-module of hyperholomorphic functions $f : \mathbb{B} \rightarrow \mathbb{H}$ satisfying

$$\sup_{a \in \mathbb{B}} \int_{\mathbb{B}} |\overline{D}f(x)|^2 (1 - |\varphi_a(x)|^2)^p dx < \infty.$$

In this talk we study several properties of these and another related spaces.

Keywords and phrases: quaternion, Bloch, Dirichlet and \mathcal{Q}_p -spaces,

Joint work with: L. M. Tovar S. (ESFM-IPN-MEX)

REFERENCES

- [1] R. Aulaskari, P. Lappan, *Criteria for an analytic function to be Bloch and a harmonic or meromorphic function to be normal*, Complex analysis and its applications, Pitman Research Notes in Math **305**, Longman Scientific and Technical Harlow (1994), 136–146.
- [2] R. Aulaskari, J. Xiao and R. Zhao, *On subspaces and subsets of BMOA and UBC*. Analysis, **15** (1995), 101–102.
- [3] K. Gürlebeck, U. Kähler, M. Shapiro, L.M. Tovar, *On \mathcal{Q}_p -Spaces of Quaternion-Valued Functions*. Complex Variables **39**, OPA N. Y. (1999) , 115–135.

(←)

CHANTAL ROTH
 Sägeweg 12, 3283 (Kallnach, Switzerland)
 (croth@nobilitas.com)

Quaternionic foundations of Quantum Mechanics and spin 1/2 visualization

Abstract. This talk explores the intriguing link between quaternion quantum mechanics and the model of the elastic continuum proposed by Augustin Cauchy, highlighting the potential of this interdisciplinary connection to advance our understanding of quantum physics. We elucidate the derivation of non- and relativistic wave equations from the Cauchy theory and Hamilton quaternion algebra, demonstrating how the quaternion representation of the elastic continuum provides a comprehensive picture of quantum mechanics. This innovative interpretation addresses the ontological challenges often associated with quantum theory, offering a means to generalize and test the theory.

Transitioning from the theoretical to the practical, we delve into a unique visualization of spinors using quaternions. We elucidate how the quaternionic representation enables a comprehension of the Dirac equation and the concept of spin 1/2, pivotal to the Planck-Kleinert model and the Pauli Exclusion Principle. We present an interactive simulation of a periodically twisting 3D grid, visualizing the spin 1/2 behavior of fermions in an elastic solid.

By visualizing Dirac spinors using two orthogonal quaternions, we illuminate the complex rotation behavior of spin 1/2 particles, drawing analogies with the famous Dirac belt trick. We define two quaternion functions that represent rotations as functions of time and distance, respectively. The visualization of the resultant spinor function as a series of concentric shells offers a illuminating representation of quantum spin, enhancing our understanding of its behavior in both time and space.

Keywords and phrases: quaternion

(←)

OLGA ROVENSKA

Institute of Mathematics of NAS of Ukraine (Kyiv, Ukraine)

(rovenskaya.olga.math@gmail.com)

Approximation of classes of Poisson integrals by Fejér means

Abstract. Asymptotic equality for upper bounds of deviations of Fejér means on classes $G(q)$ was obtained in [1] (see definitions):

$$(1) \quad \mathcal{E}(G(q); \sigma_n[f]) := \sup_{f \in G(q)} \|f(\cdot) - \sigma_n[f](\cdot)\|_C = \frac{4q}{\pi n(1+q^2)} + O(1) \frac{q^n}{n}, \quad q \in (0; q_0],$$

where $q_0 = \sqrt{2 + \sqrt{5}} - 2\sqrt{2 + \sqrt{5}} \approx 0.346$, $O(1)$ is a quantity uniformly bounded with respect to n . Also, in [1] may be found an overview of the literature. The aim of this work is to present asymptotic equality for upper bounds of deviations of Fejér means taken over classes of Poisson integrals in case $q \in [q_0; 1)$.

Theorem. Let $f \in G(q)$. For $q \in [q_0; 1)$ the equality hold as $n \rightarrow \infty$

$$(2) \quad \mathcal{E}(G(q); \sigma_n[f]) = \frac{2}{\pi n} \frac{(1+q^2)^2}{(1-q^2)(1-q^2 + \sqrt{2(1+q^4)})} + O(1) \frac{q^n}{n(1-q)^3},$$

where $O(1)$ is uniformly bounded with respect to n, q .

Keywords and phrases: Poisson integral, Fejér mean; asymptotic equality

REFERENCES

- [1] O. Novikov and O. Rovenska, Approximation of periodic analytic functions by Fejer sums, *Matematychni Studii*. **47** No 2 (2017), 196–201.

(←)

RUSLAN SALIMOV

Institute of Mathematics of the National Academy of Sciences of Ukraine (Kyiv, Ukraine)
(ruslan.salimov1@gmail.com)

MARIIA STEFANCHUK

Institute of Mathematics of the National Academy of Sciences of Ukraine (Kyiv, Ukraine)
(stefanmv43@gmail.com)

On the exponential asymptotic of solutions of the nonlinear Cauchy-Riemann-Beltrami type equation

Abstract. Let G be a domain in a complex plane \mathbb{C} , so a connected and open subset \mathbb{C} , and let $\mu: G \rightarrow \mathbb{C}$ be a measurable function with $|\mu(z)| < 1$ a.e. (almost everywhere) in G . *Beltrami equation* is called an equation of the form

$$(1) \quad f_{\bar{z}} = \mu(z)f_z,$$

where $f_{\bar{z}} = \frac{1}{2}(f_x + if_y)$, $f_z = \frac{1}{2}(f_x - if_y)$, $z = x + iy$, f_x and f_y are partial derivatives of the mapping f of x and y , respectively.

Let $\sigma: G \rightarrow \mathbb{C}$ be a measurable function and $m \geq 0$. Consider in the polar coordinate system (r, θ) the following equation:

$$(2) \quad f_r = \sigma(re^{i\theta})|f_\theta|^m f_\theta,$$

where f_r and f_θ are partial derivatives of the mapping f of r and θ , respectively. Considering the formulas $rf_r = zf_z + \bar{z}f_{\bar{z}}$, $f_\theta = i(zf_z - \bar{z}f_{\bar{z}})$, the equation (2) may be written in a complex form:

$$(3) \quad f_{\bar{z}} = \frac{z}{\bar{z}} \frac{\sigma(z)|z|}{\sigma(z)|z|} \frac{|zf_z - \bar{z}f_{\bar{z}}|^m - 1}{|zf_z - \bar{z}f_{\bar{z}}|^{m+1} + 1} f_z.$$

The mapping $f: G \rightarrow \mathbb{C}$ is called *regular at the point* $z_0 \in G$, if f has a complete differential and its Jacobian $J_f = |f_z|^2 - |f_{\bar{z}}|^2 \neq 0$ at this point. The homeomorphism f of the Sobolev class $W_{\text{loc}}^{1,1}$ is called *regular* if $J_f > 0$ a.e. A *regular homeomorphic solution* of the equation (3) is called regular homeomorphism $f: G \rightarrow \mathbb{C}$, that satisfies the equation (3) a.e. in G .

Later on we use the following notations

$$\gamma_r = \{z \in \mathbb{C} : |z| = r\}, \quad \mathbb{A}(0, r_1, r_2) = \{z \in \mathbb{C} : r_1 < |z| < r_2\}, \quad \mathbb{B} = \{z \in \mathbb{C} : |z| < 1\}.$$

Theorem. Let $m > 0$, $f: \mathbb{B} \rightarrow \mathbb{C}$ be a regular homeomorphic solution of the equation (3) of the Sobolev class $W_{\text{loc}}^{1,2}$ with the normalization $f(0) = 0$ and $\int_{\gamma_t} \frac{ds}{(\text{Im } \sigma(z))^{\frac{1}{m+1}}} \neq \infty$ for almost all $t \in (0, 1)$. Suppose that for some numbers $C > 0$, $p > 0$, $0 \leq \alpha \leq \frac{m+2}{m+1}$ and $\varepsilon_0 \in (0, 1)$ the following condition is fulfilled

$$\int_{\mathbb{A}(0, \varepsilon, \varepsilon_0)} \frac{\Phi_p^{\frac{m+2}{m+1}}(|z|) dx dy}{|z| \left(\text{Im } \overline{\sigma(z)} \right)^{\frac{1}{m+1}}} \leq C e^{\frac{\alpha}{\varepsilon^p}},$$

for any $\varepsilon \in (0, \varepsilon_0)$, where $\Phi_p(t) = \frac{e^{\frac{1}{tp}}}{t^{p+1}}$. Then

$$\liminf_{z \rightarrow 0} |f(z)| e^{\frac{\beta}{|z|^p}} \leq c_0 p^{\frac{m+2}{m}} C^{\frac{m+1}{m}} < \infty,$$

where $\beta = \frac{m+2-\alpha(m+1)}{m}$ and $c_0 = (2\pi)^{-\frac{m+1}{m}} m^{-\frac{1}{m}}$.

Keywords and phrases: Beltrami equation, regular homeomorphic solution of the Beltrami equation

REFERENCES

- [1] R. R. Salimov, M. V. Stefanchuk, *Functional asymptotic of solutions of the nonlinear Cauchy-Riemann-Beltrami system*, Nonlinear Oscillations **25** (2022), no. 4, 388–403, (in Ukrainian).

(←)

JACEK SOSNOWSKI
 NCBJ (Świerk, Poland)
 (sosnowski.jacek@wp.pl)

Influence of quantized vortex dynamics on superconducting electronics

Abstract. Superconducting materials are currently the subject of interest in modern electronic technologies, especially because they exhibit quantum phenomena on a macroscopic scale. The most evident examples are quantized vortices transporting magnetic flux, which movement determines the macroscopic current carrying phenomena in the superconducting materials, used in superconducting electronics. In the communication will be discussed the influence of the movement of the pancake type vortices in HTc multilayered superconductors in the flux creep process on their current-voltage characteristics and then critical current density, which are the essential parameters characterizing the superconducting electronic quantum devices, as well as power superconducting apparatus, as superconducting magnets with current leads. The authors works related to the superconducting cables will be reviewed shortly too. In presentation will be discussed how various initial states of the captured vortices and their shape influence the pinning potential barrier and then current-voltage characteristics. It will be presented how fitting the current-voltage characteristics to experimental data in the model allows to determine the material parameters of the HTc superconductors such as nano-sized defects concentration. Next critical current density of the superconducting materials in the vortex dynamics model will be analyzed as the function of the irradiation dose creating the nano-sized defects for various values of the magnetic induction. Initial enhancement of the critical current results from the new pinning centers creation during irradiation. At the same time, the characteristic maximum of this dependence and next decrease is interpreted as related to the destructive action on the crystal lattice of the high dose of irradiation. The vortex dynamics analysis of HTc superconductors is applied for the interpretation of the dynamic anomalies of the current-voltage characteristics in a slowly varying magnetic field, which effect can be used in superconducting electromagnetic sensors, as well as for the description of the irradiation from long Josephson's junctions with rotating vortex between the Josephson's junctions covers. Finally, the results of the previous investigations of the superconducting semiconductors with vacancies of lanthanum $\text{La}_{3-x}\text{Vac}_x\text{Se}_4$ will be discussed, possible class of the materials useful for the construction of the joined superconducting – semiconducting electronic quantum devices.

(←)

ŁUKASZ STĘPIEŃ

The Pedagogical University of Cracow (Kraków, Poland)
(sfstepie@cyf-kr.edu.pl, lukasz.stepien@up.krakow.pl)

On some exact solutions of certain nonlinear partial differential equations

Abstract. In this presentation some exact solutions (the so-called functionally-invariant solutions) of certain nonlinear partial differential equations (some heavenly equations and Landau-Lifshitz equation), will be presented. The functionally-invariant solutions were found by first time in [3]. The results presented in this talk, are obtained using the so-called structural decomposition method, which was presented and applied in [4], [5], [6], [7], [8], [9]. This method differs from the methods applied by other Authors, for e.g. [1].

Keywords and phrases: [functionally-invariant solutions, heavenly equations, Landau-Lifshitz equation]

REFERENCES

- [1] E. L. Aero, A. N. Bulygin, Y. V. Pavlov *Solutions of the sine-Gordon equation with a variable amplitude*. Theor. Math. Phys. **184**, 961–972 (2015).
- [2] M. Lakshmanan, *The fascinating world of the Landau–Lifshitz–Gilbert equation: an overview*, Phil. Trans. R. Soc. A **369**, 1280 (2011).
- [3] S. L. Sobolev, *Functionally invariant solutions of wave equation*, Travaux Inst. Physico–Math. Stekloff, Acad. Sci. **5**, 259 (1934).
- [4] Ł. T. Stępień, *Bogomolny decomposition in the context of the concept of strong necessary conditions*, PhD Thesis, Institute of Physics, Jagiellonian University, Kraków, Poland (June, 2003).
- [5] Ł. T. Stępień, *On Some Classes of Exact Solutions of Eikonal Equation*, in: "Proceedings of the 9th International Workshop on Complex Structures, Integrability and Vector Fields. Trends in Differential Geometry, Complex Analysis and Mathematical Physics", K. Sekigawa, V. S. Gerdjikov and S. Dimiev (Editors), 210 (2009).
- [6] Ł. T. Stępień, *Some decomposition method for analytic solving of certain nonlinear partial differential equations in physics with application*, J. Comp. Appl. Math. **233**, 1607 (2010).
- [7] Ł. T. Stępień, *On Some Solutions of Certain Versions of ‘Sigma’ Model and Some Skyrme-Like Models* in: "Proceedings of 9th ISAAC 2013 Congress (Kraków, Poland)", series "Current Trends in Analysis and Its Applications", V. V. Mityushev and M. V. Ruzhansky (eds.), 273 (2015).
- [8] Ł. T. Stępień, *On some exact solutions for self-dual Yang-Mills equations and generalized Yang’s equations* in: "Proceedings of 10th ISAAC 2015 Congress (Macau, China)", series "New Trends in Analysis and Interdisciplinary Applications", Pei Dang et al. (eds.), 327 (2017).
- [9] Ł. T. Stępień, *On some exact solutions of heavenly equations in four dimensions*, AIP Adv., **10**, 065105 (2020).

(←)

OSAMU SUZUKI
Nihon University (Tokyo, Japan)
(1944butterfly@gmail.com)

Binary and ternary structures in evolution theory (VI)
An entropic description of the evolution of the universe
(Open discussions)

Abstract. A method of entropy and non-commutative Galois theory for the evolution of the universe is presented. Open discussions will be expected. New ideas are wanted

Joint work with: Julian Ławrynowicz, Małgorzata Nowak-Kępczyk

REFERENCES

- [1] J. Ławrynowicz, K. Nôno, and O. Suzuki, *Binary and ternary Clifford analysis vs. non-commutative Galois extensions. I. Basics of the comparison*, Bull. Soc. Sci. Lettres Łódź Sér. Rech. Déform. **62**, no. 1 (2012), 33–42.
- [2] J. Ławrynowicz, M. Nowak-Kępczyk, O. Suzuki, in collaboration with M. Othman, *Binary and ternary structures in the evolution of the universe III. The entropy description of the evolution*, Bull. Soc. Sci. Lettres Łódź Sér. Rech. Déform. **70** no. 1, (2020) 11–41. DOI: 10.26485/0459-6854/2020/70.1/1.
- [3] Julian Ławrynowicz, Małgorzata Nowak-Kępczyk, Osamu Suzuki, *Bi-nary and ternary structures of the evolutions in the universe (2x3x2x...-world) IV. The entropy description of evolution.*, Bull. Soc. Sci. Lettres Łódź Sér. Rech. Déform. **70** no. 1, (2020) 43–81. DOI: 10.26485/0459-6854/2020/70.1/2
- [4] M. Ławrynowicz, M. Nowak-Kępczyk, O. Suzuki, *Binary and ternary structures in the evolution of the universe V, Non-commutative Galois theory of evolution*, in: (HYPER)COMPLEX Seminar 2021 In Memoriam Of Prof. Julian Ławrynowicz. A Monograph, Soc. Sci. Lodziensis, Łódź 2021, pp. 7–37. DOI: 10.26485/978-83-60655-92-4/2

(←)

VASILEIOS VACHTSEVANOS
Aristotle University of Thessaloniki (Thessaloniki, Greece)
(vvachtse@physics.auth.gr)

Emergence of topological defects in finite 2-D lattices during second phase order transitions

Abstract. In this work we study the Kibble-Zurek [1] mechanism that describes the emergence of topological defects in finite 2-D lattices during second phase order transitions. We will discuss the numerical method used to recreate the process, mention the behaviour of 1-D atomic chains and expound on the difficulty behind the categorization of the 3 different types of defects (point, 1-D and 2-D) when using statistical methods. We will shortly discuss the mathematics concerning both the nature of the defects and the numerical process (Euler-Maruyama Leapfrog).

Our work is based on previous work that is done concerning atomic chains [2]-[5].

Joint work with: Hariton Polatoglou

REFERENCES

- [1] Zurek, Wojciech H, 'Cosmological experiments in condensed matter systems'upload, Physics Reports 276 (1996), (4), 177-221.
- [2] Vachtsevanos V. Polatoglou H., 'Statistics of topological defects in finite one-dimensional structures based on the Kibble Zurek Mechanism', Journal of Vibration Testing and System Dynamics, Accepted 16/04/2023.
- [3] Del Campo, Adolfo.'Universal statistics of topological defects formed in a quantum phase transition', Physical review letters 121 (2018), (20), 200601.
- [4] Del Campo, A.; Kibble, T. W. B.; Zurek, W. H., 'Causality and non-equilibrium second-order phase transitions in inhomogeneous systems', Journal of Physics: Condensed Matter, 25 (2013) (40), 404210.
- [5] Gomez-Ruiz, Fernando J.; Mayo Jack J.; Del Campo, Adolfo., 'Full counting statistics of topological defects after crossing a phase transition', Physical Review Letters 124 (2020) (24), 240602.

(←)

VALERY VOLCHKOV
 Donetsk National University (Donetsk, Ukraine)
 (volna936@gmail.com)

VITALY VOLCHKOV
 Donetsk National University (Donetsk, Ukraine)
 (volna936@gmail.com)

Continuous mean periodic extension of functions from a segment

Abstract. Let $n \geq 2$, $\mathcal{E}'(\mathbb{R}^n)$ be the space of compactly supported distributions on \mathbb{R}^n . We study the following version of the mean periodic extension problem.

(i) Suppose that $T \in \mathcal{E}'(\mathbb{R}^n)$, and E is a non-empty closed subset of \mathbb{R}^n . What conditions guarantee that for a function $f \in C(E)$ there is a function $F \in C(\mathbb{R}^n)$ coinciding with f on E and satisfying the convolution equation $F * T = 0$ in \mathbb{R}^n ?

(ii) If such an extension F exists then estimate the growth of F at infinity.

We present a solution of this problem for a broad class of distributions T in the case when E is a segment in \mathbb{R}^n .

For results related to the continuation of mean periodic functions, see [1]-[3] and the references there.

Keywords and phrases: Convolution equations, mean periodicity, spherical transform, quasi-analyticity

REFERENCES

- [1] V. V. Volchkov, *Integral Geometry and Convolution Equations*, Dordrecht: Kluwer Academic Publishers, 2003, (ISBN 1-4020-1628-X).
- [2] V. V. Volchkov, Vit. V. Volchkov, *Harmonic Analysis of Mean Periodic Functions on Symmetric Spaces and the Heisenberg Group*, London: Springer-Verlag, 2009, (ISBN 978-1-84882-532-1).
- [3] V. V. Volchkov, Vit. V. Volchkov, *Offbeat Integral Geometry on Symmetric Spaces*, Basel: Birkhäuser, 2013, (ISBN 978-3-0348-0571-1).

(←)

OLEH YERMAKOV

Department of Computer Physics, V. N. Karazin Kharkiv National University (Kharkiv, Ukraine)
(oe.yermakov@gmail.com)

SERGEY POLEVOY

O. Ya. Usikov Institute of Radiophysics and Electronics NASU (Kharkiv, Ukraine)

Discovery of polarization degree of freedom for localized light

Abstract. The spectrum of electromagnetic plane waves in any isotropic homogeneous medium is twice degenerate with respect to polarization at any frequency and in any direction, because the dispersions of TE- and TM-polarized eigenmodes are absolutely the same. The operational principle of any bulk classical polarizer is based on the removal of the polarization degeneracy by using an anisotropic slab.

Miniaturization and planar technologies lead to the high localization of the electromagnetic signal in the plane of propagation. However, at the same time, the degeneracy is removed. The simplest example is the dielectric slab waveguide demonstrating the propagation of highly localized guided modes with different dispersions of TE and TM modes. So, there is no polarization degree of freedom for planar photonic devices with in-plane electromagnetic wave propagation, since the TE and TM localized modes are not degenerate. It significantly limits the functionality of flat optical and planar photonic devices. In particular, the polarizer of guided waves cannot be implemented without the broadband TE-TM polarization degeneracy.

In this work, we show the possible resolution of this fundamental problem via the nanostructuring of the dielectric slab. We aim merging the polarization degeneracy and high localization for guided waves propagating within an all-dielectric metasurface. Namely, we theoretically study the periodic subwavelength array of cylinders with high refractive index and demonstrate the broadband TE-TM polarization degeneracy of its guided modes. This concept can be transferred to the optical and infrared ranges using other material platforms and opens new opportunities for the polarization control of guided waves such as metasurface-based waveguide polarizer [1].

Another possible implementation includes the self-complementary metasurfaces obeying Babinet's duality principle [2]. In this case, we excite independently the TE- and TM-polarized surface waves at self-complementary metasurfaces. Then, by adjusting the complex amplitudes of TE and TM eigenmodes, we demonstrate the excitation of surface waves of linear, circular and arbitrary elliptical polarization [3].

These results could potentially become a platform for new generation of planar photonic polarization devices.

Keywords and phrases: photonics, polarization, metasurfaces

REFERENCES

- [1] O. Y. Yermakov, A. A. Bogdanov, A. V. Lavrinenko, *Broadband polarization degeneracy of guided waves in subwavelength structured ZnO pattern*, IEEE Journal of Selected Topics in Quantum Electronics **25**, 3 (2019).
- [2] O. Yermakov, V. Lenets, A. Sayanskiy, J. Baena, E. Martini, S. Glybovski, S. Maci, *Surface waves on selfcomplementary metasurfaces: All-frequency hyperbolicity, extreme canalization, and TE-TM polarization degeneracy*, Physical Review X **11**, 031038 (2021).
- [3] S. Polevoy, O. Yermakov, *Excitation of surface waves with on-demand polarization at self-complementary metasurface*, IEEE Antennas and Wireless Propagation Letters, Early Access (2023).

(←)

MIKOŁAJ ZŁOTKOWSKI
AGH, Faculty of Energy and Fuels (Kraków, Poland)
(mikoazolot@interia.pl)

Local systems management of different energy sources

Abstract. The question arises: Is the correct solution for energy information system having container controlling heat storage, electric energy sources that are unstable or diverse in time? The goal is to obtain high-quality energy from OZE as from energetic system grid or burning methane or other fuels. The key to success is set as the miniaturization of consumption energy for device AGD. Is it necessary to turn down things in the house that shouldn't be on power all day? The presented method is the first step in distributed energy networks (RSE), adaptation to smart house technology, and management strategy for controlling of nodes of energy supply system. In such a way a lot of houses in village/city have creative cluster storage energy system. This solution reduces the load on PSE. Parallel to PSE working local direct power lines, alternative lines are used to control flow energy from OZE to storage devices. The presentation gives not very new concepts, but provides a specific diagram of the operation of a high-power system about 1MW.

(←)

MARIUSZ ZUBERT

Lodz University of Technology, Department of Microelectronics and Computer (Lodz, Poland)
(mariusz.zubert@p.lodz.pl)

Application of Scattering Parameters to The DPL Time-Lag Parameter Estimation using MEMS Structures

Abstract. The dedicated micro-electro-mechanical system (MEMS) test structure has been developed and fabricated for material parameters' estimation for the dual-phase-lag (DPL) model at the nanoscale in modern integration circuit (IC) structures [1, 2]. The temperature time-lag estimation procedure of dielectrics is presented based on the electric scattering parameters measured by a vector network analyser for the considered MEMS test structure together with the 2-omega method [2]. The proposed methodology has the ability to estimate the time-lag parameter with high accuracy and is also suitable for the temperature time-lag estimation for other manufacturing process technologies of ICs for insulation materials such as silicon dioxide (SiO₂), silicon nitride (Si₃N₄), titanium nitride (TiN), and hafnium dioxide (HfO₂).

Keywords and phrases: dual-phase-lag heat transfer model; no-mesh FDM; S-parameters measurements; scattering parameters; DPL time lag estimation; multi-domain modelling; material parameter estimation; heat transfer at the nanoscale

Joint work with: Z. Kulesza, M. Jankowski, A. Napieralski

REFERENCES

- [1] M. Zubert M, T. Raszkowski, A. Samson, P. Zajac, *Methodology of determining the applicability range of the DPL model to heat transfer in modern integrated circuits comprised of FinFETs*. *Microelectronics Reliability*, vol. **91**, 2018, 139—153, (<https://doi.org/10.1016/j.microrel.2018.07.141>).
- [2] M. Zubert, Z. Kulesza, M. Jankowski, A. Napieralski, *A. Application of Scattering Parameters to DPL Time-Lag Parameter Estimation at Nanoscale in Modern Integration Circuit Structures*. *Energies* 2021, **14**, 4425. (<https://doi.org/10.3390/en14154425>)

(←)

NATALIA ZORII

Institute of Mathematics of National Academy of Sciences of Ukraine (Kyiv, Ukraine)
(natalia.zorii@gmail.com)

**On the role of the point at infinity in Deny's principle of positivity of mass
for Riesz potentials**

Abstract. This talk is based on [9], and it deals with the theory of potentials on \mathbb{R}^n , $n \geq 2$, with respect to the Riesz kernel $|x - y|^{\alpha - n}$, $\alpha \in (0, 2]$, $\alpha < n$, where $|x - y|$ is the Euclidean distance between $x, y \in \mathbb{R}^n$. Denote by \mathfrak{M}^+ the cone of all positive Radon measures μ on \mathbb{R}^n such that the Riesz potential

$$U^\mu(x) := \int |x - y|^{\alpha - n} d\mu(y)$$

is not identically infinite on \mathbb{R}^n , which according to [5, Section I.3.7] occurs if and only if

$$\int_{|y|>1} \frac{d\mu(y)}{|y|^{n-\alpha}} < \infty.$$

Then U^μ is actually finite everywhere on \mathbb{R}^n , up to a set of *zero Riesz capacity*, cf. [5, Section III.1.1].

The principle of positivity of mass was first introduced by J. Deny (see e.g. [2]), and for Riesz potentials it reads as follows [3, Theorem 3.11].

Theorem 1. *For any $\mu, \nu \in \mathfrak{M}^+$ such that*

$$(1) \quad U^\mu \leq U^\nu \quad \text{everywhere on } \mathbb{R}^n,$$

we have $\mu(\mathbb{R}^n) \leq \nu(\mathbb{R}^n)$.

It is easy to verify that (1) can be slightly weakened by replacing ‘everywhere on \mathbb{R}^n ’ by ‘nearly everywhere on \mathbb{R}^n ’ (see [8, Theorem 2.6], establishing the principle of positivity of mass for potentials with respect to rather general function kernels on locally compact spaces). Recall that a proposition $\mathcal{P}(x)$ is said to hold *nearly everywhere (n.e.)* on $A \subset \mathbb{R}^n$ if $c_*(E) = 0$, where E is the set of all $x \in A$ for which $\mathcal{P}(x)$ fails, while $c_*(E)$ denotes the *inner Riesz capacity* of E , see [5, Section II.2.6].

The main result of this talk, given by Theorem 2, shows that Theorem 1 still holds even if (1) is fulfilled only on a proper subset A of \mathbb{R}^n , which however must be ‘large enough’ in an arbitrarily small neighborhood of $\infty_{\mathbb{R}^n}$, the Alexandroff point of \mathbb{R}^n . This discovery illustrates a special role of the point at infinity in Riesz potential theory, in particular with regard to the principle of positivity of mass.

Theorem 2. *Given $\mu, \nu \in \mathfrak{M}^+$, assume there exists $A \subset \mathbb{R}^n$ which is not inner α -thin at infinity, and such that*

$$U^\mu \leq U^\nu \quad \text{n.e. on } A.$$

Then

$$\mu(\mathbb{R}^n) \leq \nu(\mathbb{R}^n).$$

Recall that according to [4, 7], $A \subset \mathbb{R}^n$ is said to be *inner α -thin at infinity* if

$$\sum_{k \in \mathbb{N}} \frac{c_*(A_k)}{q^{k(n-\alpha)}} < \infty,$$

where $q \in (1, \infty)$ and $A_k := A \cap \{x \in \mathbb{R}^n : q^k \leq |x| < q^{k+1}\}$; or equivalently, if either A is bounded, or $x = 0$ is an inner α -irregular boundary point for the inverse of A with respect to $|x| = 1$. (For the concept of inner α -irregular points for arbitrary $A \subset \mathbb{R}^n$ and relevant results, see [6, Section 6]; compare with [5, Section V.1], where A was required to be Borel.) We emphasize that *if A is not inner α -thin at infinity, then necessarily $c_*(A) = \infty$; but not the other way around* (see [7, Section 2]).

The following theorem shows that Theorem 2 is *sharp* in the sense that the requirement on A of not being α -thin at infinity can not in general be weakened.

Theorem 3. *If $A \subset \mathbb{R}^n$ is inner α -thin at infinity, then there are $\mu_0, \nu_0 \in \mathfrak{M}^+$ such that $U^{\mu_0} = U^{\nu_0}$ nearly everywhere on A , but nonetheless, $\mu_0(\mathbb{R}^n) > \nu_0(\mathbb{R}^n)$.*

Nevertheless, Theorem 2 remains valid for *arbitrary* $A \subset \mathbb{R}^n$ once we impose upon $\mu, \nu \in \mathfrak{M}^+$ suitable additional requirements (see Theorem 4 below).

A measure $\mu \in \mathfrak{M}^+$ is said to be *carried* by $A \subset \mathbb{R}^n$ if $\mathbb{R}^n \setminus A$ is μ -negligible, or equivalently if A is μ -measurable and $\mu = \mu|_A$, $\mu|_A$ being the trace of μ to A , cf. [1, Section V.5.7]. We denote by \mathfrak{M}_A^+ the cone of all $\mu \in \mathfrak{M}^+$ carried by A . (For closed A , μ is carried by A if and only if it is supported by A .)

A measure $\mu \in \mathfrak{M}^+$ is said to be *C-absolutely continuous* if $\mu(K) = 0$ for every compact set $K \subset \mathbb{R}^n$ of zero Riesz capacity. This certainly occurs if $\int U^\mu d\mu$ is finite (or, more generally, if U^μ is locally bounded); but not conversely, see [5, pp. 134–135].

Theorem 4. *For any set $A \subset \mathbb{R}^n$ and any C-absolutely continuous measures $\mu, \nu \in \mathfrak{M}_A^+$ such that $U^\mu \leq U^\nu$ n.e. on A , we still have $\mu(\mathbb{R}^n) \leq \nu(\mathbb{R}^n)$.*

Remark 1. If $A \cap A_I = \emptyset$, where A_I denotes the set of all inner α -irregular points for A , then the requirement of C-absolute continuity imposed on μ and ν , is unnecessary for the validity of Theorem 4.

Remark 2. The proofs of the above-quoted theorems are based on the theory of inner α -Riesz balayage as well as on that of inner α -Riesz equilibrium measures, both originated in [6, 7] (see also [8]). The concept of inner equilibrium measure is understood in an extended sense where its energy as well as its total mass may be infinite. The following two facts of these theories are crucial to our proofs:

- $A \subset \mathbb{R}^n$ is not α -thin at infinity if and only if the inner balayage of *any* $\mu \in \mathfrak{M}^+$ to A preserves its total mass (see [7, Corollary 5.3]).
- The inner α -Riesz equilibrium measure of $A \subset \mathbb{R}^n$ exists if and only if A is α -thin at infinity (see [7, Theorem 2.1]).

Remark 3. The results presented in the talk have already found applications to minimum Riesz energy problems in the presence of external fields, see for instance [10, Section 4.10].

REFERENCES

- [1] Bourbaki, N.: *Integration*. Chapters 1–6. Springer, Berlin (2004)
- [2] Deny, J.: *Méthodes Hilbertiennes en Théorie du Potentiel*, In: *Potential Theory*. CIME. Summer Schools 49, pp. 121–201. Springer, Berlin (2010)
- [3] Fuglede, B., Zorii, N.: *Green kernels associated with Riesz kernels*, *Ann. Acad. Sci. Fenn. Math.* **43**, 121–145 (2018)
- [4] Kurokawa, T., Mizuta, Y.: *On the order at infinity of Riesz potentials*, *Hiroshima Math. J.* **9**, 533–545 (1979)
- [5] Landkof, N.S.: *Foundations of Modern Potential Theory*, Springer, Berlin (1972)
- [6] Zorii, N.: *A theory of inner Riesz balayage and its applications*, *Bull. Pol. Acad. Sci. Math.* **68**, 41–67 (2020)
- [7] Zorii, N.: *Harmonic measure, equilibrium measure, and thinness at infinity in the theory of Riesz potentials*, *Potential Anal.* **57**, 447–472 (2022)
- [8] Zorii, N.: *On the theory of capacities on locally compact spaces and its interaction with the theory of balayage*, *Potential Anal.* (2022). <https://doi.org/10.1007/s11118-022-10010-3>
- [9] Zorii, N.: *On the role of the point at infinity in Deny’s principle of positivity of mass for Riesz potentials*, *Anal. Math. Phys.* **13**, 38 (2023)
- [10] Zorii, N.: *Minimum Riesz energy problems with external fields*, *J. Math. Anal. Appl.* **526**, 127235 (2023)

(←)